

Photon Structure in γp Interactions^{ab}

J. H. VOSSEBELD

NIKHEF, P.O. Box 41882, 1009 DB Amsterdam, The Netherlands

E-mail: vossebeld@nikhef.nl

On behalf of the ZEUS collaboration.

Photoproduction of jets at HERA provides information on the partonic structure of the photon. We report on the latest dijet photoproduction results, for real photons and for photons at low virtualities, measured with the ZEUS detector.

1 Introduction

In the ep -collider HERA^c we can study, in addition to ep Deep Inelastic Scattering, also γp scattering. The photoproduction of jets is a process, well suited to test perturbative QCD and to study the partonic structure of the photon and the proton. In this paper we concentrate on the structure of the photon.

In leading order we distinguish two classes of photoproduction processes: in the *direct* process the photon couples directly to a parton in the proton while in the *resolved* process a parton inside the photon couples to a parton in the proton. Kinematically the two classes can be separated using x_γ , the fractional momentum of the photon participating in the hard scatter. For direct events $x_\gamma \sim 1$ while for resolved events $x_\gamma < 1$. In resolved processes the partonic structure of the photon is probed.

We study events in which at least two hard jets are produced. Hard jets are associated with high P_T partons produced in the subprocess. The observation of hard jets thus ensures the presence of a hard scale. We present cross section measurements using two types of jetfinders. An iterative cone algorithm, according to the Snowmass convention², and a k_T -clustering algorithm³. For comparison of data with higher order QCD calculations the k_T -clustering algorithm is preferred. The “Snowmass” cone algorithm is not unambiguously defined⁴ and has numerical instabilities at higher orders⁵.

2 Dijet photoproduction

Dijet photoproduction is particularly well suited to study the partonic structure of the photon, because the reconstruction of two jets in the final state allows us

^aTo appear in the proceedings of the DIS98 workshop held in Brussels (Belgium), April 4-8, 1998.

^bThe inclusive jet photoproduction results presented at the workshop have been published¹, and will not be covered in this paper

^cFor the data presented in this paper 820 GeV protons were collided with 27.5 GeV positrons.

to determine the kinematics of the subprocess. From the transverse energy (E_T^{jet}) and the pseudorapidity^d (η^{jet}) of the jets one obtains the fractional momentum of the photon participating in the hard scatter:

$$x_\gamma^{obs} = \left(E_{T1}^{jet} e^{-\eta_1^{jet}} + E_{T2}^{jet} e^{-\eta_2^{jet}} \right) / 2yE_e \quad (1)$$

where E_e is the beam energy of the incoming positron and y is the fractional energy transfer from the positron to the proton in the proton rest frame.

The dijet cross section is measured as a function of the transverse energy and the pseudorapidity of the jets in the following kinematic range: $-1 < \eta_{1,2}^{jet} < 2$, $E_{T1}^{jet} > 14 \text{ GeV}$, $E_{T2}^{jet} > 11 \text{ GeV}$, $0.20 < y < 0.85$ and $Q^2 < 1 \text{ GeV}^2$. Jets are defined using a k_T -clustering algorithm.

From previous jet photoproduction analyses at lower transverse energies^{1,7}, we have learned that the cross sections show an excess over Monte Carlo predictions for forward jets, or correspondingly, for low x_γ^{obs} values. The data are well described when a multiple interaction model is included in the Monte Carlo simulation.




Figure 1: The uncorrected x_γ^{obs} spectrum (left plot) and the transverse energy flow around jets (right plots), compared to Herwig 5.9. The transverse energy is plotted as a function of the distance in pseudorapidity with respect to the jet (integrated over $|\Delta\phi| < 1$).

Requiring two hard jets reduces the sensitivity of the measured cross section to multiple interaction effects. As is shown in figure 1 the measured spectra of x_γ^{obs} and the transverse energy flow around jets are in excellent agreement with predictions of the Herwig 5.9⁸ Monte Carlo, which has been used without including the multiple interaction option.

Figure 2 and 3 show the dijet cross section as a function of the transverse energy of the jets and of the pseudorapidity of the jets. Note that in both cases the cross section is symmetrized with respect to η_1^{jet} and η_2^{jet} , therefore every

^d $\eta^{jet} \equiv -\ln(\tan \frac{\theta^{jet}}{2})$

^eThe application of an asymmetric cut on the transverse energies of the two jets is required to ensure the stability of the NLO calculations⁶.




Figure 2: Inclusive dijet cross section as a function of E_T^{jet} for various combinations of η_1^{jet} and η_2^{jet} . The gray band indicates the (strongly correlated) uncertainty related to the energy scale.




Figure 3: Inclusive dijet cross section as a function of η_2^{jet} while η_1^{jet} is fixed. The upper (lower) curves correspond to the full (high) x_γ^{obs} range. The full (dashed) curves are Kramer and Klasen NLO calculations using the GS96 (GRV) parameterization for the photon structure. The dotted curves are Harris and Owens calculations using GS96 for the photon structure.

event enters twice, once for every jet. Both for the full x_γ^{obs} -range and for the high x_γ^{obs} -range ($x_\gamma^{obs} > 0.75$), the data agree well with NLO QCD predictions⁹. We note that the differences related to the used photon parameterization¹⁰ are of similar size as the uncertainty in the measurement.

The dijet cross section is also measured as a function of the pseudorapidities of the jets in a restricted kinematic range ($0.50 < y < 0.85$) to increase the sensitivity to the photon structure. Effectively this restriction removes the highest x_γ (direct) events in every bin. In figure 4 the data are compared to NLO calculations.




Figure 4: Inclusive dijet cross section as a function of η_2^{jet} while η_1^{jet} is fixed, for high values of y . The full (dashed) curves correspond to Klasen and Kramer NLO calculations using the GS96 (GRV) parameterization for the photon structure.

tions using the same parameterizations of the photon structure as in figure 3. An increased sensitivity to the photon structure is observed in the NLO calculations. At central pseudorapidities, a significant discrepancy between the data and the QCD predictions is observed.

Scale uncertainties⁹ and hadronization effects are expected to be below 10%.

3 Dijet production at low photon virtualities

A subject of great interest is the production of dijets when the photon has a finite virtuality Q^2 . The contribution from resolved processes to the cross section is expected to be reduced for increasing values of Q^2 .

We have selected a dijet sample in the following kinematic range: $-1.125 < \eta_{1,2}^{jet} < 1.875$, $E_{T1,2}^{jet} > 6.5 \text{ GeV}$ and $0.2 < y < 0.55$. Jets are defined with a cone algorithm.

The dijet cross section is determined in two ranges of Q^2 : the untagged photo-production region for which $Q^2 \sim 0$ and the range $0.10 < Q^2 < 0.70 \text{ GeV}^2$, which corresponds to events in which the scattered positron is detected in the beam pipe calorimeter¹¹.




Figure 5: The uncorrected x_γ^{obs} spectrum (left plot) in different bins of Q^2 , compared to Herwig 5.9 (no multiple interactions and no Q^2 suppression). The hatched histogram shows the direct component of the Monte Carlo separately. The right plot shows the “resolved over direct” ratio as a function of Q^2 .

In figure 5 the x_γ^{obs} distribution is compared to Herwig 5.9, in different bins of Q^2 . For Q^2 values up to 0.70 GeV^2 , both the resolved and the direct (dark histogram) component of the Monte Carlo are needed to describe the data.

Also in figure 5 the “resolved over direct” ratio of the cross section for $x_\gamma^{obs} < 0.75$ over that for $x_\gamma^{obs} > 0.75$ is plotted as a function of Q^2 . A decrease of

this ratio with Q^2 is observed. Herwig 5.9, which contains no mechanism for Q^2 suppression of the resolved component, can not reproduce this behavior. NLO calculations¹² using the (Q^2 suppressed) SaS¹³ parameterization for the photon, also fail to describe the data. This might be related to the symmetric cut applied to the transverse energy of the two jets (see footnote e).

4 Summary

Photoproduction of jets at HERA offers a unique source of information on the structure of the photon. The dijet photoproduction measurements presented in this paper show a discrepancy between data and NLO QCD predictions for high energy real photons. For virtual photons we have observed a decrease of the “resolved over direct” ratio as a function of Q^2 (up to $Q^2 = 0.70 \text{ GeV}^2$).

References

1. J.Breitweg et al, *DESY* 98-018, accepted by *Europ. Phys. Jour.*
2. F. Abe et al, *Phys. Rev. D* **45**, 1448 (1992)
3. S. Catani, Yu.L. Dokshitzer, M.H. Seymour and B.R. Webber, *Nucl. Phys. B* **406**, 187 (1993);
S.D. Ellis and D.E. Soper, *Phys. Rev. D* **48**, 3160 (1993)
4. S.D. Ellis, Z. Kunszt and D.E. Soper, *Phys. Rev. Lett.* **69**, 3615 (1992)
5. M.H. Seymour, *HEP-PH/9707349*
6. M. Klasen and G. Kramer, *Phys. Lett. B* **366**, 385 (1996);
S. Frixione and G. Ridolfi, *Nucl. Phys. B* **507**, 315 (1997)
7. J.Breitweg et al, *Europ. Phys. J. C* **1**, 109 (1998);
S. Aid et al, *Z. Phys. C* **70**, 17 (1996);
M.Derrick et al, *Phys. Lett. B* **348**, 665 (1995)
8. G. Marchesini, B.R. Webber, G. Abbiendi, I.G. Knowles, M.H. Seymour and L. Stanco, *Comput. Phys. Commun.* **67**, 465 (1992);
G. Marchesini, B.R. Webber, G. Abbiendi, I.G. Knowles, M.H. Seymour and L. Stanco, *HEP-PH/9607393*
9. M. Klasen, T. Kleinwort and G. Kramer, *DESY* 97-234;
B.W. Harris and J.F. Owens, *Phys. Rev. D* **57**, 5555 (1998)
10. M. Glück, E. Reya and A. Vogt, *Phys. Rev. D* **46**, 1973 (1992);
M. Glück, E. Reya and A. Vogt, *Phys. Rev. D* **45**, 3986 (1992);
L.E. Gordon and J.K. Storrow, *Nucl. Phys. B* **489**, 405 (1997)
11. B. Surrow, PhD thesis Univ. Hamburg, Jan. 1998
12. M. Klasen, G. Kramer and B. Pötter, *Europ. Phys. J. C* **1**, 261 (1998)
13. G.A. Schuler and T. Sjöstrand, *Phys. Lett. B* **376**, 193 (1996)